



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Methodological bias associated with soluble protein recovery from soil

Citation for published version:

Greenfield, LM, Hill, PW, Paterson, E, Baggs, EM & Jones, DL 2018, 'Methodological bias associated with soluble protein recovery from soil', *Scientific Reports*, vol. 8, no. 1, pp. 11186.
<https://doi.org/10.1038/s41598-018-29559-4>

Digital Object Identifier (DOI):

[10.1038/s41598-018-29559-4](https://doi.org/10.1038/s41598-018-29559-4)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Scientific Reports

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



SCIENTIFIC REPORTS

OPEN

Methodological bias associated with soluble protein recovery from soil

Lucy M. Greenfield¹ , Paul W. Hill¹, Eric Paterson², Elizabeth M. Baggs³ & Davey L. Jones¹

Proteins play a crucial role in many soil processes, however, standardised methods to extract soluble protein from soil are lacking. The aim of this study was to compare the ability of different extractants to quantify the recovery of soluble proteins from three soil types (Cambisol, Ferralsol and Histosol) with contrasting clay and organic matter contents. Known amounts of plant-derived ¹⁴C-labelled soluble proteins were incubated with soil and then extracted with solutions of contrasting pH, concentration and polarity. Protein recovery proved highly solvent and soil dependent (Histosol > Cambisol > Ferralsol) and no single extractant was capable of complete protein recovery. In comparison to deionised water (10–60% of the total protein recovered), maximal recovery was observed with NaOH (0.1 M; 61–80%) and Na-pyrophosphate (0.05 M, pH 7.0; 45–75% recovery). We conclude that the dependence of protein recovery on both extractant and soil type prevents direct comparison of studies using different recovery methods, particularly if no extraction controls are used. We present recommendations for a standard protein extraction protocol.

Protein represents the dominant form of organic nitrogen (N) entering soil ecosystems and frequently the bottleneck in soil N cycling¹. Further, based on the number of proteins contained in plants and microorganisms, it can be expected that a single gram of soil may contain thousands of different proteins^{2,3}. As proteins play a key role in many soil processes, there is increasing interest in the extraction, separation, identification and quantification of proteins as indicators of soil function. However, the development of exoproteomic approaches are currently limited by the lack of standard protocols and the difficulty of recovering proteins from soil.

Extractants that have commonly been used for soil protein recovery include simple salts (e.g. K₂SO₄, Na-pyrophosphate, Na-phosphate), bases (e.g. NaOH), organic acids (e.g. Na-citrate) and surfactants (e.g. Tris-SDS) (Supplementary Table S1). Although previous studies have examined a range of protein extraction methods, these have been largely restricted to single unrepresentative proteins (e.g. BSA), single soils or have used quantification methods known to suffer from severe interference by the co-extraction of humic substances^{4–6}. In addition, many of these studies have lacked the appropriate controls, preventing determination of protein extraction efficiency or have focused on the whole soil metaproteome.

Soil type has a large influence on protein recovery. Some studies suggest that organic matter and clay content are the key soil properties which affect protein recovery^{5,7,8} whilst other studies suggest soil pH is also important^{9–11}. Organic matter content, clay content and pH influence the adsorption of protein in soil and, therefore, affects the ease to which it can be extracted.

Our aim was to focus on soluble proteins and to compare the recovery of a mixture of ¹⁴C-labelled plant proteins from soil using 39 different extractants. Our secondary aim was to evaluate the influence of soil type on protein recovery.

Materials and Methods

Soils used in the study. We evaluated protein recovery from three soils with contrasting organic matter and Fe contents: (1) a Eutric Cambisol obtained from a temperate *Lolium perenne* L. grassland in Abergwyngregyn, Gwynedd, UK (53°14'N, 4°00'W); (2) a Fibric Histosol obtained from a temperate *Calluna vulgaris* (L.) Hull moorland in Abergwyngregyn, Gwynedd, UK (53°22'N, 4°01'W), and (3) a Rhodic Ferralsol obtained from a

¹School of Environment, Natural Resources & Geography, Bangor University, Gwynedd, LL57 2UW, UK. ²The James Hutton Institute, Craigiebuckler, Aberdeen, AB15 8QH, UK. ³The Royal (Dick) School of Veterinary Studies, University of Edinburgh, Easter Bush Campus, Midlothian, EH25 9RG, UK. Correspondence and requests for materials should be addressed to L.M.G. (email: l.greenfield@bangor.ac.uk)

| | Cambisol | Ferralsol | Histosol |
|---|--------------------------|---------------------------|--------------------------|
| pH | 6.07 ± 0.02 ^a | 4.75 ± 0.22 ^b | 4.48 ± 0.17 ^b |
| EC (μS cm ⁻¹) | 21.8 ± 3.3 ^a | 140.7 ± 52.9 ^a | 47.5 ± 28.4 ^a |
| Organic C (%) | 2.43 ± 0.02 ^a | 1.45 ± 0.08 ^a | 23.2 ± 0.5 ^b |
| Total N (%) | 0.21 ± 0.00 ^a | 0.12 ± 0.00 ^b | 1.12 ± 0.04 ^c |
| Sand (%) | 40.7 ± 3.2 ^a | 31.0 ± 1.8 ^a | 90.7 ± 2.4 ^b |
| Silt (%) | 46.0 ± 2.7 ^a | 35.1 ± 2.2 ^b | 8.0 ± 1.9 ^c |
| Clay (%) | 13.3 ± 0.5 ^a | 33.8 ± 0.7 ^b | 1.3 ± 0.5 ^c |
| Cation exchange capacity (mmol kg ⁻¹) | 145 ± 6 ^a | 90 ± 8 ^a | 334 ± 57 ^b |

Table 1. Major characteristics of the three soils used in the extraction trial. Values represent means ± SEM (Standard error of the mean) ($n = 3$). Different letters (^a, ^b, ^c) indicate significant differences between soils at the $p < 0.05$ level.

Saccharum officinarum L. plantation in Piracicaba, Brazil (22°32'S, 49°20'W)¹². In all cases, replicate batches of soil ($n = 3$) were collected from a depth of 0–15 cm, sieved (<2 mm) and kept at 4 °C until required. The main soil properties are shown in Table 1. Soil pH and electrical conductivity (EC) were measured in 1:5 (v/v) soil:H₂O extracts. Total C and N were determined with a TruSpec[®] analyser (Leco Corp., St Joseph, MI). Soil texture was determined with a LS1330 Particle size analyser (Beckman Coulter, Brea, CA). Cation exchange capacity (CEC) was measured by saturation with an index cation¹³. Soluble protein in water extracts was measured using the Coomassie Blue method¹⁴ and was used to calibrate the rate of ¹⁴C-labelled protein addition (Supplementary Table S3). This method, however, cannot be used with other extractants other than water due to bias from interfering substances^{6,15}.

Protein extraction solutions. The extractants tested were based on previously published methods (Supplementary Table S1) and included: deionised water, Na-pyrophosphate (0.01, 0.05, 0.1 M; pH 7.0), Na-citrate (0.01, 0.05, 0.1, 0.5 M; pH 8.0), Tris-SDS (0.01, 0.05, 0.1 M SDS with 0.05 M Tris; pH 7.0), K-phosphate buffer (0.01, 0.05, 0.1, 0.5 M; pH 6.0 and 8.0), CaCl₂, NaOH and K₂SO₄ (0.01, 0.05, 0.1, 0.5 M), methanol and ethanol (25%, 50%, 75%, 100% v/v). Extractants with no pH value stated were not adjusted and their values are presented in Supplementary Table S4.

Protein addition and recovery from soil. Soil (1 g) was placed in individual 20 ml polypropylene vials and heat-sterilised (80 °C, 1 h) immediately prior to experimentation¹⁶. This sterilisation procedure was not found to affect the CEC of the soils (Supplementary Table S2). In addition, it also proved effective at killing the microbial community preventing bias from microbial breakdown/immobilisation of the added protein (Supplementary Fig. S1). Although free protease activity was not completely eliminated by heat sterilisation, the exoenzyme activity was extremely low compared to the amount of protein added to the soil and was therefore not expected to bias our findings (Supplementary Table S5). Purified, ¹⁴C-uniformly labelled soluble protein from *Nicotiana tabacum* L. leaves (100 μl; 0.860 mg ml⁻¹; 1.2 kBq ml⁻¹; purified to >3 kDa by ultra-filtration; custom synthesised by American Radiolabeled Chemicals, St Louis, MO) was added to each soil, shaken to mix and incubated for 30 min at 20 °C. An incubation time of 30 min was deemed appropriate based on initial pilot studies of protein sorption and recovery from soil at incubation times varying from 0.5 to 24 h (Supplementary Table S6). The time is therefore sufficient to obtain high rates of sorption while minimising the chances of proteolysis or microbial regrowth. Soluble plant proteins were chosen as they represent one of the major forms of dissolved organic N added to soil. Based on extractant methods from previous studies, the soils were subsequently shaken with 5 ml of each extractant (30 min; 200 rev min⁻¹)^{17,18}, then a 1.5 ml aliquot was pipetted into 1.5 ml microfuge tubes and centrifuged (18 000 g; 60 s) and the supernatant recovered. The centrifugation time of 60 s allowed complete phase separation of the soil particles and supernatant (Supplementary Table S7). The amount of ¹⁴C-label recovered in degradations per minute (DPM) of supernatant was determined using a Wallac 1414 scintillation counter (60 s) and Wallac Optiphase HiSafe3 scintillation fluid (PerkinElmer Inc., Waltham, MA). Baseline ¹⁴C-labelled protein was determined by counting 100 μl of ¹⁴C-labelled protein. Extraction efficiency was calculated by Equation (1).

$$\text{Extraction efficiency (\%)} = \frac{{}^{14}\text{C in extractant supernatant (DPM)}}{\text{baseline } {}^{14}\text{C (DPM)}} \times 100 \quad (1)$$

Humic acids and organic solvents had no effect on ¹⁴C counting efficiency (Supplementary Tables S8 and S9). To estimate the amount of humic substances co-extracted with the protein, the colour of the extracts was determined at 254 and 400 nm in UV-transparent plastic 96-well plate using a PowerWave HT Spectrophotometer (BioTek Inc., Winooski, VT).

Statistical analysis. All experiments were performed in triplicate. All statistical analysis was performed using R 3.4.1 and work was carried out in base R unless stated¹⁹. Data was declared to be normally distributed by Shapiro-Wilk normality test ($p > 0.05$) and have equal variances across groups by Bartlett test ($p > 0.05$). Graphs were created using the R package ggplot2²⁰. Differences in soil properties between soil types were analysed by one-way ANOVA with TukeyHSD post-hoc testing using $p < 0.05$ as the cut-off for statistical significance.




Figure 1. Extraction efficiency (%) of ^{14}C -labelled protein from three contrasting soils using a range of chemical extractants. The legend to the left of the dashed line refers concentrations of methanol and ethanol. For all other extractants, refer to the legend on the right of the dashed line. Different capital letters represent significant differences between soil type of the same molarity and extractant. Different letters represent significant differences between molarity of the same soil type and extractant. Values represent means \pm SEM ($n = 3$).

Differences in protein recovery between treatments and soils were analysed by two-way ANOVA with TukeyHSD post-hoc testing using $p < 0.05$ as the cut-off for statistical significance. Chemical speciation modelling to estimate the net valency of each extractant was performed with Geochem-EZ²¹.

Data availability. Please contact the corresponding author (l.greenfield@bangor.ac.uk) for access to data.

Results and Discussion

Protein recovery from soil by water. Here we aimed to evaluate methods of soluble protein recovery. This is relevant to studies investigating the potential behaviour of isotopically labelled proteins in soil (sorption, biodegradation) or for recovering the plant or microbial exoproteome. Overall, we found significant differences in protein extraction efficiency between the different extractants ($F_{10, 318} = 118.5$; $p < 0.001$; Fig. 1) and soils (Histosol > Cambisol > Ferralsol) ($F_{2, 318} = 148.4$; $p < 0.001$; Fig. 1). As the soil was sterilised to limit microbial activity¹⁶, ^{14}C measured is assumed to represent intact ^{14}C -protein, therefore we refer to extraction efficiency as protein recovered. Protein recovery by deionised water varied from 10–60% between soil types. As the water is expected to recover mainly free, unbound protein, we assume the remainder became bound to the solid phase or coagulated/precipitated on entering the soil^{22,23}. Proteins are known to readily sorb to the surface of clay minerals, Fe/Al oxyhydroxides and humic materials in soil^{24–26}. Therefore, extractants should be able to displace proteins bound to surfaces during the extraction process or to solubilise the binding surfaces themselves. Ferralsols had the lowest protein recovery probably because of the higher clay and Fe-oxide fraction, compared to the Histosol and Cambisol ($p < 0.001$; Table 1), resulting in more protein being strongly bound to the solid phase. In comparison, the higher humic content of the Histosol ($p < 0.001$; Table 1) may have resulted in the extraction of protein as soluble humic-protein complexes²⁷. In our soils, a complete recovery of the added ^{14}C -protein was not achieved for any soil, with ca. 25% not recoverable by any extractant. This is likely to be even higher in soils where proteins have been stabilised for long periods.

Protein recovery from soil by salt extracts. For the Histosol, no significant difference was observed between deionised water and the other extractants ($p > 0.05$) except CaCl_2 and K_2SO_4 which lowered protein extraction compared to deionised water ($p < 0.05$). We ascribe the poor protein recovery with CaCl_2 and K_2SO_4 to salt-induced conformational changes in protein structure and subsequent coagulation/precipitation (Supplementary Table S10), a phenomenon which is well documented in the literature²³. In contrast to the

Histosol, deionised water gave low protein recovery rates from the Ferralsol and Cambisol likely due to more protein adsorbed onto the clay fraction. We conclude therefore that water extracts may provide an estimate of free, unbound proteins in soil and limited information of the bound fraction. Further, while 0.5 M K_2SO_4 is frequently used as a standard extractant for dissolved organic N and for measuring soil microbial biomass-N^{17,28}, our results suggest that the method may reduce total protein recovery.

The highest recoveries were obtained by NaOH and Na-pyrophosphate (70–76% of the total protein added), with no significant difference apparent between them ($p > 0.05$; Fig. 1). The high pH of NaOH relative to the other extractants solubilises organic matter leading to the release of protein particularly the case of the Histosol²⁹. For the Ferralsol, NaOH was the most efficient extractant (49–77% compared to 43–48% by Na-citrate). NaOH also solubilises protein adsorbed to $Al(OH)_3$, resulting in protein release from the Ferralsol³⁰.

Our results therefore suggest that the recovery of protein from soil is consistent with (i) their salting-out potential based on the Hofmeister series²³, and (ii) the potential of each salt to displace bound protein from surfaces via ligand exchange, based on their net valency (i.e. $HP_2O_7^{3-} > Citrate^{3-} > phosphate^{1.87- (pH 8)} = phosphate^{1.15- (pH 6)} > SO_4^{2-} > Cl^-$). The exception to this was Tris-SDS^{0.09} which had a significantly higher extraction efficiency than K_2SO_4 and $CaCl_2$ ($p < 0.001$) suggesting that the presence of surfactant aids ionic displacement. Surfactants tend to gather around interfaces (e.g. the interface between the soil surface and soil solution). The surfactants compete with the protein molecules for available surface area in order for the hydrophobic tails to avoid water. Over time the SDS molecule will replace the protein molecules because the surfactant molecules are in excess^{31,32}.

Protein recovery from soil by organic solvents. The polar solvents, methanol and ethanol both proved ineffectual at recovering soluble proteins from soil likely due to the alcohol-induced precipitation of proteins³³. This contrasts strongly with metabolomic studies where these extractants often yield the greatest recovery of low molecular weight organic solutes^{34,35}.

Co-extraction of humic substances. NaOH caused the solubilisation of large amounts of humic substances and based on previous studies, this is likely to induce protein denaturation^{36,37}. Consequently, we would not recommend it as an extractant. However, in some analysis the structure of the protein is not important (e.g. sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and the Kjeldahl method) and NaOH can be used.

Na-pyrophosphate, NaOH, Na-citrate and both phosphate buffers extracted more humic substances in comparison to deionised water (Supplementary Table S11; Supplementary Fig. S2) in support of previous findings^{11,38}. Humic substances can be problematic due to their ability to bind to proteins and interfere with colorimetric procedures for quantifying protein^{4,39}. Proteins interact with humic substances to form protein-humic substance complexes²⁶. The mechanisms of the interaction are thought to consist of: (a) covalent and hydrogen bonds⁴⁰, (b) ionic bonds between the functional amino group of the protein and the carboxyl or hydroxyl group of the humic substance⁴¹, (c) physically immobilised within macromolecular matrix of humic substances⁴², and (d) electron donor-acceptor complexes⁴³.

The co-extraction of humic substances with proteins in the protein-humic complexes results in colour in the supernatant. This interferes with colorimetric and fluorescent analysis of protein quantity^{6,44}. Methods of removing interfering humic substances (e.g. PVPP⁴ and TCA precipitation⁴⁵) have been found to be ineffective⁴⁶. Therefore, NaOH, Na-pyrophosphate, Na-citrate and phosphate buffers are not ideal extractants when these types of analysis are being used. In addition, if extracting protein from a soil with high organic matter content, more interference will occur in comparison to soils with lower organic matter contents.

Conclusions

In summary, we found that 0.1 M NaOH was the most effective extractant overall when denatured protein can be used in subsequent analysis and co-extraction of humic substances does not interfere. For analysis of intact proteins, 0.05 M Na-pyrophosphate (pH 7.0) was most effective for extracting water-soluble proteins from soil; however, it did also co-extract humic substances. Where interference of humic substances may prove problematic for subsequent analysis and intact proteins are required, deionised water is recommended. For proteomics, further analysis by LC-MS/MS will be necessary to assess the quality of the proteins extracted by each method^{15,47}. In addition, although this study was limited to three soils, our results clearly indicate that soil type directly affects the amount of protein that can be recovered. This may make quantitative comparisons between soils problematic. Rarely has this been accounted for in previous studies comparing protein levels in soil. The impact of this in future studies can be evaluated by measuring the recovery of a known mixture of proteins, as undertaken here. It should also be emphasised that this study focused only on the recovery of hydrophilic proteins from soil. Similar studies are therefore required to optimise the recovery of proteins contained within the soil microbial community, especially those of a hydrophobic nature.

References

1. Jan, M. T., Roberts, P., Tonheim, S. K. & Jones, D. L. Protein breakdown represents a major bottleneck in nitrogen cycling in grassland soils. *Soil Biol. Biochem.* **41**, 2272–2282 (2009).
2. Haas, B. J. *et al.* Complete reannotation of the *Arabidopsis* genome: methods, tools, protocols and the final release. *BMC Biol.* **3**, 1–19 (2005).
3. Ishihama, Y. *et al.* Protein abundance profiling of the *Escherichia coli* cytosol. *BMC Genomics* **9**, 1–19 (2008).
4. Criquet, S., Farnet, A. M. & Ferre, E. Protein measurement in forest litter. *Biol. Fertil. Soils* **35**, 307–313 (2002).
5. Kanerva, S., Smolander, A., Kitunen, V., Ketola, R. A. & Kotiaho, T. Comparison of extractants and applicability of MALDI-TOF-MS in the analysis of soil proteinaceous material from different types of soil. *Org. Geochem.* **56**, 1–9 (2013).
6. Roberts, P. & Jones, D. L. Critical evaluation of methods for determining total protein in soil solution. *Soil Biol. Biochem.* **40**, 1485–1495 (2008).

7. Bastida, F., Jehmlich, N., Torres, I. F. & García, C. The extracellular metaproteome of soils under semiarid climate: A methodological comparison of extraction buffers. *Sci. Total Environ.* **619–620**, 707–711 (2018).
8. Chen, S., Rillig, M. C. & Wang, W. Improving soil protein extraction for metaproteome analysis and glomalin-related soil protein detection. *Proteomics* **9**, 4970–4973 (2009).
9. Haney, R. L., Franzluebbers, A. J., Hons, F. M., Hossner, L. R. & Zuberer, D. A. Molar concentration of K_2SO_4 and soil pH affect estimation of extractable C with chloroform fumigation-extraction. *Soil Biol. Biochem.* **33**, 1501–1507 (2001).
10. Halvorson, J. J. & Gonzalez, J. M. Bradford reactive soil protein in Appalachian soils: distribution and response to incubation, extraction reagent and tannins. *Plant Soil* **286**, 339–356 (2006).
11. Masciandaro, G. *et al.* Comparison of extraction methods for recovery of extracellular β -glucosidase in two different forest soils. *Soil Biol. Biochem.* **40**, 2156–2161 (2008).
12. IUSS Working Group WRB. *World reference base for soil resources 2014, update 2015 International soil classification system for naming soils and creating legends for soil maps. World Soil Resources Reports No. 106* (2015).
13. Rhoades, J. D. In *Methods of soil analysis. Part 2. Chemical and Microbiological Properties* (eds Page, A. L., Miller, R. H. & Keeney, D. R.) 149–157 (Soil Science Society of America, 1982).
14. Bradford, M. M. A Rapid and Sensitive Method for the Quantitation of Microgram Quantities of Protein Utilizing the Principle of Protein-Dye Binding. *Anal. Biochem.* **72**, 248–254 (1976).
15. Bastida, F., Hernández, T. & García, C. Metaproteomics of soils from semiarid environment: Functional and phylogenetic information obtained with different protein extraction methods. *J. Proteomics* **101**, 31–42 (2014).
16. Mariano, E., Jones, D. L., Hill, P. W. & Trivelin, P. C. O. Mineralisation and sorption of dissolved organic nitrogen compounds in litter and soil from sugarcane fields. *Soil Biol. Biochem.* **103**, 522–532 (2016).
17. Jones, D. & Willett, V. Experimental evaluation of methods to quantify dissolved organic nitrogen (DON) and dissolved organic carbon (DOC) in soil. *Soil Biol. Biochem.* **38**, 991–999 (2006).
18. Rousk, J. & Jones, D. L. Loss of low molecular weight dissolved organic carbon (DOC) and nitrogen (DON) in H_2O and 0.5 M K_2SO_4 soil extracts. *Soil Biol. Biochem.* **42**, 2331–2335 (2010).
19. R Core Team. R: A language and environment for statistical computing (2017).
20. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis* (2016).
21. Shaff, J. E., Schultz, B. A., Craft, E. J., Clark, R. T. & Kochian, L. V. GEOCHEM-EZ: a chemical speciation program with greater power and flexibility. *Plant Soil* **330**, 207–214 (2010).
22. Coen, C. J., Blanch, H. W. & Prausnitz, J. M. Salting out of aqueous proteins: Phase equilibria and intermolecular potentials. *Am. Inst. Chem. Eng.* **41**, 996–1004 (1995).
23. Shih, Y.-C., Prausnitz, J. M. & Blanch, H. W. Some characteristics of protein precipitation by salts. *Biotechnol. Bioeng.* **40**, 1155–1164 (1992).
24. Quiquampoix, H. *et al.* In *Proteins at Interfaces II. Fundamentals and applications* (eds Horbett, T. & Brash, J. L.) 321–333 (American Chemical Society, 1995).
25. Yu, W. H. *et al.* Adsorption of proteins and nucleic acids on clay minerals and their interactions: A review. *Appl. Clay Sci.* **80**, 443–452 (2013).
26. Nielsen, K. M., Calamai, L. & Pietramellara, G. In *Nucleic Acids and Proteins in Soil* (eds Nannipieri, P. & Smalla, K.) 141–157 (Springer-Verlag, 2006).
27. Bonmati, M., Ceccanti, B., Nannipieri, P. & Valero, J. Characterization of humus–protease complexes extracted from soil. *Soil Biol. Biochem.* **41**, 1199–1209 (2009).
28. Joergensen, R. G. & Brookes, P. C. Ninhydrin-reactive nitrogen measurements of microbial biomass in 0.5 M K_2SO_4 soil extracts. *Soil Biol. Biochem.* **22**, 1023–1027 (1990).
29. Schnitzer, M. In *Methods of Soil Analysis. Part 2. Chemical and Microbiological Properties* (eds Page, A. L., Miller, R. H. & Keeney, D. R.) 581–594 (American Society of Agronomy, Soil Science Society of America, 1982). doi:10.2134/agronmonogr9.2.2ed.c30.
30. Gianfreda, L., Rao, M. A. & Violante, A. Adsorption, activity and kinetic properties of urease on montmorillonite, aluminium hydroxide and $Al(OH)_3$ -montmorillonite complexes. *Soil Biol. Biochem.* **24**, 51–58 (1992).
31. Norde, W. My voyage of discovery to proteins in flatland ... and beyond. *Colloids Surfaces B Biointerfaces* **61**, 1–9 (2008).
32. Jönsson, A.-S. & Jönsson, B. The influence of nonionic and ionic surfactants on hydrophobic and hydrophilic ultrafiltration membranes. *J. Memb. Sci.* **56**, 49–76 (1991).
33. Schubert, P. F. & Finn, R. K. Alcohol precipitation of proteins: The relationship of denaturation and precipitation for catalase. *Biotechnol. Bioeng.* **23**, 2569–2590 (1981).
34. Swenson, T. L., Jenkins, S., Bowen, B. P. & Northen, T. R. Untargeted soil metabolomics methods for analysis of extractable organic matter. *Soil Biol. Biochem.* **80**, 189–198 (2015).
35. Warren, C. R. Comparison of methods for extraction of organic N monomers from soil microbial biomass. *Soil Biol. Biochem.* **81**, 67–76 (2015).
36. Bremner, J. M. In *Soil Biochemistry* (eds McLaren, A. D. & Peterson, G. H.) 19–58 (Marcel Dekker Inc., 1967).
37. Bremner, J. M. & Lees, H. Studies on soil organic matter: Part II. The extraction of organic matter from soil by neutral reagents. *J. Agric. Sci.* **39**, 274–279 (1949).
38. Murase, A., Yoneda, M., Ueno, R. & Yonebayashi, K. Isolation of extracellular protein from greenhouse soil. *Soil Biol. Biochem.* **35**, 733–736 (2003).
39. Bastida, F., Moreno, J. L., Nicolás, C., Hernández, T. & García, C. Soil metaproteomics: A review of an emerging environmental science. *Significance, methodology and perspectives. Eur. J. Soil Sci.* **60**, 845–859 (2009).
40. Rowell, M. J., Ladd, J. N. & Paul, A. A. Enzymically active complexes of proteases and humic acid analogues. *Soil Biol. Biochem.* **5**, 699–703 (1973).
41. Sarkar, J. M. Formation of $[^{14}C]$ cellulase-humic complexes and their stability in soil. *Soil Biol. Biochem.* **18**, 251–254 (1986).
42. Serban, A. & Nissenbaum, A. Humic acid association with peroxidase and catalase. *Soil Biol. Biochem.* **18**, 41–44 (1986).
43. Gosewinkel, U. & Broadbent, F. E. Decomplexation of phosphatase from extracted soil humic substances with electron donating reagents. *Soil Sci.* **141**, 261–267 (1986).
44. Whiffen, L. K., Midgley, D. J. & McGee, P. A. Polyphenolic compounds interfere with quantification of protein in soil extracts using the Bradford method. *Soil Biol. Biochem.* **39**, 691–694 (2007).
45. Qian, C. & Hettich, R. L. Optimized Extraction Method To Remove Humic Acid Interferences from Soil Samples Prior to Microbial Proteome Measurements. *J. Proteome Res.* **16**, 2537–2546 (2017).
46. Aoyama, M. Properties of neutral phosphate buffer extractable organic matter in soils revealed using size exclusion chromatography and fractionation with polyvinylpyrrolidone. *Soil Sci. Plant Nutr.* **52**, 378–386 (2006).
47. Leary, D. H., Hervey, W. J., Deschamps, J. R., Kusterbeck, A. W. & Vora, G. J. Which metaproteome? The impact of protein extraction bias on metaproteomic analyses. *Mol. Cell. Probes* **27**, 193–199 (2013).

Acknowledgements

This work was supported by the Biotechnology and Biological Sciences Research Council; and the Natural Environment Research Council [Grant number NE/M009106/1], by a Soils Training and Research Studentships (STARS) grant to LMG. STARS is a consortium consisting of Bangor University, British Geological Survey, Centre for Ecology and Hydrology, Cranfield University, James Hutton Institute, Lancaster University, Rothamsted Research and the University of Nottingham.

Author Contributions

L.G., D.J. and P.H. conceived the experiment. L.G. conducted the experiment and wrote the main manuscript. L.G., D.J., P.H., E.P. and E.B. reviewed the manuscript.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-018-29559-4>.

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018